
Remarks on bryophyte collections from N-Spain including the first record of *Dicranodontium didymodon* in Europe

Jan-Peter Frahm

Abstract: *Fontinalis antipyretica* var. *mollis* is recorded for Spain, which differs by orbicular, not keeled leaves. The presence in the same tufts together with var. *antipyretica* reveals that it is a somatic mutant of the latter. A form of *Pellia epiphylla* was observed which has wide thallus margins of unistratose cells resembling those of *Aneura maxima*. A *Campylopus* was collected resembling *C. flexuosus* but with ventral stereids in transverse section of the costa. It is described as *C. flexuosus* var. *anomalus*. An (unsuccessful) attempt has been made to find the locality in which P. Allorge found *Campylopus setifolius* for the only time in the Iberian Peninsula. The bryophilous fungus *Octospora jungermanniarum* is reported as new to Spain. *Dicranodontium didymodon*, known before from the Himalaya and western China, is reported as new to Europe.

From February 27 to March 7, 2013, I made a fieldtrip through northern Spain (Cantabria, Asturias, Galicia) in order to study the atlantic bryoflora. An enumeration of the collections is omitted here, since the distribution of most of the species seems to be well known, however, some otherwise remarkable collections as well as notes on previous records are compiled here.

A somatic mutant of *Fontinalis antipyretica*

Whilst collecting aquatic mosses (*Cinclidotus fontinaloides*, *C. riparius*, *Fontinalis antipyretica*) in the Rio Puron (near the village of Puron in the vicinity of Llanes, -4,700197, 43,385116, alt. 27 m), a strange *Fontinalis* was found with almost orbicular, very concave leaves and no indication of a keel. Such a *Fontinalis* was already reported from Germany, where it was collected in the river Sülz near Cologne (Frahm 2009). Except for *Fontinalis cymbifolia* Nichols., no other species of the genus from Europe matches this specimen, however, a *F. antipyretica* var. *mollis* (C. Müll.) Welch in Grout was described earlier from North America and therefore this name was tentatively applied. Interestingly var. *mollis* was described as a variety of *F. antipyretica*, although the round leaves without keel would suggest that the taxon would belong even to a different subgenus. In Germany, *F. antipyretica* var. *mollis* was associated in the river with normal expressions of *F. antipyretica*, which proved that var. *mollis* is genetically distinct and no modification when

growing together with the type variety. This seemed to be also the case in Spain until single stems of the var. *mollis* were found growing intermingled in tufts of var. *antipyretica*. This indicates that the *mollis* expression is a somatic mutant of var. *antipyretica*.

The shape of the leaves of var. *mollis* resembles that of the perichaetial leaves and it seems as if a genetical switch is thrown to produce perichaetial leaves along a normal stem.

Cases of somatic mutants are known and genetically proved from other mosses such as *Hypnum cupressiforme* and *Platyhypnidium riparioides*. The mutant of *H. cupressiforme* with julaceous foliation was described as *H. heseleri* and that of *P. riparioides* with multitratose lamina as *P. mutatum*. In the case of *Hypnum*, single branches of the *heseleri* expression were found (van Zanten & Hofman 1994) and in *P. mutatum*, single plants were found growing in tufts of *P. riparioides* (Frahm 2005), although both species grow usually in separate tufts. *Thamnobryum fernandezii*, again an aquatic moss with multitratose lamina, is growing intermingled with *Th. alopecurum* (pers. obs.)

In all these cases the mutant was described as separate species because of the morphological and anatomical differences, although isozyme analyses (*H. heseleri*) or DNA sequences (*Platyhypnidium mutatum*, *Palustriella pluristratosa*, *Ochyrea tatrensis* etc.) revealed no or no significant differences as compared with the original species. If this use is applied to the case of *Fontinalis*, the mutant has to be named *F. mollis* C. Müll. in Röll.



Fig. 1: Branches of *Fontinalis antipyretica* var. *antipyretica* (top) and var. *mollis* (bottom) in the same tuft.

***Pellia epiphylla* with wide unistratose thallus margin**

Searching for atlantic species such as *Drepanolejeunea hamatifolia* or *Harpalejeunea molleri* in a ravine near San Sebastian de Garabandal in the valley of Rio Vendula (Cantabria, 43°42'08.4"N, 4°43'20.5882"E, alt. 420m) a thallose liverwort was found which was



Fig. 2: Thallus of *Pellia epiphylla* with broad unistratose border.



Fig. 3: The same thallus in translucent light.



Fig. 4: For comparison thallus of “normal” *Pellia epiphylla*.

as *Aneura maxima* (interestingly a synonym of *Aneura maxima* in Japan is *Aneura pellioides*). Later sporophytes were found, which showed involucre proving that the plants belong to *Pellia epiphylla*.

Principally *Pellia epiphylla* has unistratose thallus margins, which are usually rather narrow (cf. fig. 4), but in this case the broad unicellular wings of *Pellia epiphylla* (fig. 3) remind at *Aneura maxima*, for which the *Pellia* was taken at first in the field. It might be that there is a gene in the genome of both species, (or generally in the Aneuraceae and Pelliaceae), which is usually switched off in *Pellia*.

***Campylopus flexuosus* with ventral stereids**

In April 2011, Felipe Gutiérrez Pérez collected a species of *Campylopus* in NW-Spain, which had ventral stereids in transverse section of the costa. Since *Campylopus brevipilus* is the only species in Europe with ventral stereids, the specimen was tentatively attributed to this species, however, as he stated, the leaves had no hairpoints and were like in *C. flexuosus* in shape, but the basal cells near de costa porose.

Worldwide, there are several *Campylopus* species resembling *C. flexuosus* but with ventral stereids, *Campylopus arctocarpus* in the Neotropics, *C. arctocarpus* ssp. *madegassus* in tropical Africa, *C. flagelliferus* in SE-Asia, *C. cubensis* in the Carribean and *C. incorralis* in the tropical mountains. All species are presumably of common origin and geographical vicariants. For a final decision, the specimen was sequenced by Michael Stech, specialist for molecular systematics of Dicranaceae, (University of Leiden), however the marker studied placed the specimen to *C. flexuosus*.

Ventral stereids in the costa of *Campylopus flexuosus* were already reported by Frahm (1977) with regard to the infrageneric classification of the genus by Limpricht (1890) based upon the transverse section of the costa. Limpricht grouped the species in those having ventral and dorsal stereids (*Palinocraspis*), dorsal stereids (*Campylopus*) and no stereids (*Pseudocampylopus*). The fact that *Campylopus flexuosus*, the type species of the genus and subgenus *Campylopus* can have ventral hyalocysts as well as stereids, corroborate this differentiation. Recent molecular systematic studies (Stech 1999) did not maintain this classification.

The reason for the formation of ventral stereids in *C. flexuosus* is not clear. Frahm (1977) excluded different habitats or age of the three specimens mentioned from Germany.

During the author's trip, a *Campylopus* cf. *flexuosus* with ventral stereids (fig. 5,6) was collected again in Asturias, in the valley of Rio de Nueva. A new molecular study based on this material is in progress.

The problem involved is that the anatomical structure of the costa is distinctly different from *C. flexuosus*, although the sequence of the marker used is identical, and that plants with different anatomical structure deserve to be recognized as a different taxon. Therefore this aberrant expression is described here as a new variety:

Campylopus flexuosus (Hedw.) Brid. Mant. Musc. 4:71, 1819 var. *anomalus* J.-P. Frahm var. nov. Differs from the typical variety by the presence of distinct ventral stereids in the transverse section of the costa instead of ventral hyalocysts.

Type: Spain, Asturias, Rio de Nueva S of Nueva, on a N-exposed slope (known for the occurrence of *Woodwardia radicans*), 4,949374° W, 43,430688°N, alt. 200m., ca. 2 km upstream of road AS 340, leg. Frahm 2.3.2013 no. SP-046 (hb. Frahm BONN).





Fig. 5,6: Transverse section of the costa of *Campylopus* cf. *flexuosus* with ventral stereids. (Frahm SP-046)

***Campylopus setifolius* in Spain**

In 1933, Pierre Allorge collected *Campylopus setifolius* in northern Spain, new to the Iberian Peninsula (Allorge 1934) which is so far the only record. It is cited as follows:

Asturias: vallee du Rio Ore Carroyas, quartzites suintants, a 1 km environ de l'Ocean'

I have seen the specimen, which is deposited in the Natural History Museum in Paris. The specimen is labelled as follows:

rochers quartzites, valle du rio Ore, Carroyas, entre Luarca et Canero, leg. P. Allorge 12.9.33 teste Dixon.

In 2013, Gordon Rothero asked in the bryonet whether this is still the only record outside Britain and where this locality is situated, since Google failed to locate the place.

Except for some suggestions, the Spanish bryologist Montserrat Brugués gave the information:

Caroyas (47°32'44,23"N/6°28'40,72"W), río Orio (or Esba, name of this river when is near the ocean).

Allorge's information seems to be quite distinct (about 1 km near the ocean). Caroyas is in fact spelled with one r, not two. It is situated 1 kilometre away from a river which is today called rio Esva.

The supplementary information on the label (between Luarca and Canero) reveals that the valley of rio Ore must be the present rio Esva. A search for the locality in March 2013 revealed the following:

Inhabitants of the village of Caroyas do not know Rio Ore, however, in his publication, Allorge (1934) mentions more collections (such as *Saccogyna viticulosa*, *Plagiochila spinulosa* and *Bartramia stricta*) from Vallée du rio Ore with the addition "entre Castaneda et Ponteijon". This is misleading, because the village of Castaneda is south of Santander. However, there is a village of Castanedo upstream rio Esva. Downstream is a village called Pontigon, which is probably the

same as Allorge's Ponteijon. Interestingly, a village called Ore is closeby, an indication that the name of the river was rio Ore in the past.

Therefore it can be concluded that Allorge's rio Ore is the present rio Esva and the locality of *Campylopus setifolius* is 1 km south of the mouth of the river. There are, however, no natural rocks or cliffs (fig. 7). In a longer distance, an old railway bridge is crossing the valley. Here are rocks along the eastern side of the road, on which *Bryum alpinum*, *Amphidium mougeotii* on seeping rocks as well as *Campylopus introflexus*, *Targionia hypophylla* and *Trichostomum brachydontium* in drier situation are growing. Compared with the habitats in Scotland the situation here is fairly dry. It must be supposed, that the roadside rocks were cut when the road was expanded. As shown by cut offs of an old narrow road, the road was renewed, probably after Allorge's visit. Under this precondition, a suitable habitat has been destroyed, although is almost two kilometres from the sea. Due to the lack of any other suitable habitats in the surroundings, *C. setifolius* is most probably no more present in the Iberian Peninsula.

The only record of an atlantic bryophyte species in Northern Spain is nothing exceptional. Even *Lepidozia cupressina* (one of the commonest representatives of this element) or *Lejeunea mandonii* have been found there only once. Other species, for example, *Breutelia chrysocoma*, and *Isoetecium holtii*, which are frequent in areas where the species occurs, are surprisingly rare and scattered in northern Spain. This indicates that these records are probably no relics of former larger extensions but new colonizations.




Fig. 7. Google map of the mouth of Rio Esva, where Allorge found *Campylopus setifolius* in 1 km distance from the ocean (yellow line = 1 km).



Fig. 8: *Octosporella jungermanniarum* growing on *Plagiochila punctata*.

***Octosporella jungermanniarum*, a hepaticolous fungus new to Spain**

Whilst searching for *Tetralophozia filiformis* at the place where E. Urmi found this species for the first time in Europe (Asturias, 2,5 km S of Sellano at Rio Ponga, -5,211722°W, 43,232059°N, about 340 m alt.), I collected *Plagiochila punctata*. Under the dissection microscope I observed fruit bodies of a jungus (fig. 8), which was identified by Peter Döbbeler, specialist for bryophilous fungi, as *Octosporella jungermanniarum* (P. Crouan & H. Crouan) Döbbeler (Ascomycota, Pyronemataceae). According to Döbbeler, this species is new to Spain although it has been recorded about 50 times from Central Europe.

***Dicranodontium didymodon* (Griff.) Par new to Europe**

When bryologizing in the valley of Rio de Nueva¹, I collected a dicranaceous moss which looked in the field like a *Campylopus* or, what I supposed, perhaps *Dicranodontium uncinatum*, however with less uncinata leaves. A microscopic examination revealed that it was in fact a *Dicranodontium*, because of the elongate upper laminal cells, the extremely long and fine subula and the ventral stereids in transverse section of the costa. It was, however, none of the known species from Europe. These can easily be divided in two groups: species with large, lax, hyaline inner basal laminal cells (like in *Calymperes*) including *D. asperulum* and *D. uncinatum* and a

¹ Asturias, Rio de Nueva S of Nueva, on a N-exposed slope, 4,949374° W, 43,430688°N, alt. 200m., ca. 2 km upstream of road AS 340, leg. Frahm 2.3.2013 no. SP-047 (hb. Frahm BONN).

species with “normal” inner basal laminal cells such as *D. denudatum* and *D. subporodictyon*. The present plants had “normal” inner basal laminal cells, but differed from *D. denudatum* by a only finely serrate to almost smooth subula and the costa well differentiated from the lamina at the base of the leaf. In contrast, *D. denudatum* has a sharply serrate subula (like a rats tail file) and the costa is not clearly separated from the lamina at the base of the leaf. In addition, the leaves of *D. denudatum* are easily falling off or are easily detached with a wet finger, leaving a stem with rests of leaf bases which look like an old defoliate branch of a spruce.

Years ago, I had monographed *Dicranodontium* (Frahm 1997) and using the key, I ended with *D. didymodon* (Griff.) Par. This species is known from the Himalaya region, from India (Darjeeling, Sikkim), Bhutan, Nepal, northern Thailand and Burma to China (Sichuan).

With regard to this range, the occurrence in northern Spain seems to be surprising, however, *Dicranodontium asperulum*, *D. subporodictyon* and *D. uncinatum* have the same disjunction as well as numerous species, which are found in the Himalaya, Japan, western North America, the atlantic coastal regions of Europe and the alpine mountains of Europe (or parts of it). Examples are except for the three species of *Dicranodontium* mentioned above *Campylopus gracilis*, *Leptodontium recurvifolium* and especially liverworts such as *Herbertus sendtneri*, *Anastrophyllum assimile*, *A. donnianum*, *A. joergensenii*, *Pleurozia purpurea*, *Matsigophora woodsii*, *Bazzania perasonii*, *Douinia ovata*, *Anastrepta orcadensis*, *Plagiochila carringtonii*, *Scapania ornithopodioides* and *S. nimbosea*. So this new record is just a new example.

Interestingly, the liverwort *Tetralophozia filiformis* was found as new to Europe in 1978 in Asturias in the valley of Rio Ponga (Urmi 1983). It has a similar somehow wider distribution and is known from Sikkim, Darjeeling, Yunnan, Sichuan, Japan, Taiwan, Borneo and British Columbia. Later search of several bryologists could not confirm the existence at this locality anymore, however, this species found later again at more localities (e.g. Fuertes Lasala 1987) at much higher altitudes. Therefore the first locality can be regarded as untypical for this montane species.

In the valley of Rio de Nueva, *Dicranodontium* grew in tufts of *Leucobryum glaucum* on humid siliceous rocks on the base of a N-exposed slope (fig. 9), which is known for the occurrence of the fern *Woodwardia radicans*. The slope is mainly covered by *Sphagnum capillifolium*, *S. denticulatum*, *S. palustre*, *Hylocomium splendens*. Other associates were *Fissidens polyphyllus*, *Plagiochila exigua*, *P. punctata*, *Kurzia sylvatica*, *Scapania nemorea*, *Plagiothecium undulatum* and *Calypogeia neesiana*.

Dicranodontium didymodon may be overlooked in northern Spain because of its very inconspicuous appearance like a *Dicranum scoparium* or *Campylopus* (fig. 10). It is differentiated in the field with a hand lens from the first by the wide costa and from the latter only under the microscope by the characters mentioned above (fig. 11). A characteristic of *Dicranodontium* seems to be that the ventral stereids in transverse section of the costa are confined to the middle of the costa, whereas they extend over the whole costa in species of *Campylopus* with ventral stereids.

Acknowledgements

I wish to thank Marta Infante and Patxi Heras (Vitoria) for valuable suggestions of bryologically interesting localities, and Peter Döbbeler for the identification of *Octospora jungermanniarum*.



Fig. 9: Habitat of *Dicranodontium didymodon* at the Woodwardia locality in the valley of Rio de Nueva.



Fig. 10: *Dicranodontium didymodon*




Fig. 4. *Dicranodontium didymodon* (Griff.) Par. (Griffith 32, H-BR). — a: leaf. — b: Leaf base. — c: Basal laminal cells. — d: Upper laminal cells. — e: Transverse section of costa.

Fig. 11: Illustration of *Dicranodontium didymodon* from Frahm (1997).

Literature

- Allorge, P. 1934. Notes sur la flore bryologique de la Peninsule Iberique IX. Muscinées des Provinces du Nord et du Centre de l'Espagne. *Revue Bryologique* n.s. 7: 249-301.
- Frahm, J.-P. 1997. A taxonomic revision of *Dicranodontium* (Musci). *Ann. Bot. fennici* 34: 179-204.
- Frahm, J.-P. 2005. *Platyhypnidium mutatum* Ochyra & Vanderpoorten an seinem weltweit einzigen Vorkommen im Schwarzwald. *Bryol. Rundbriefe* 93: 1-2.
- Frahm, J.-P. 2009. *Fontinalis antipyretica* var. *mollis* in Deutschland. *Archive for Bryology* 59, 5 pp.
- Frahm, J.-P. 1977. Ventrale stereidale Bildungen im Rippenquerschnitt von *Campylopus flexuosus* (Hedw.) Brid. *Herzogia* 4: 249-264.
- Fuertes Lasala, E. 1987. A new locality for *Tetralophozia filiformis* (Steph.) Urmi in Spain. *J. Bryol.* 14: 791.
- Limpricht, K.G. 1890. Die Laubmoose Deutschlands, Österreichs und der Schweiz. Rabenhorsts Kryptogamenflora Bd. 4, Leipzig.
- Stech, M. 1999. A reclassification of Dicranaceae (Bryopsida) based on non-coding cpDNA sequence data. *Journal of the Hattori Botanical Laboratory* 86: 137-159.
- Urmi, E. 1983. *Tetralophozia filiformis* (Steph.) comb. nov. in Europe. *J. Bryol.* 12: 393-401.
- van Zanten, B.O., Hofman, A. 1994. On the possible origin and taxonomic status of *Hypnum heseleri* Ando & Deguchi. *J. Hattori Bot. Lab.* 75: 107-118.

Online March 22, 2013